

# Treatment Discussion Guide

To help you prepare to talk to your doctor

If your **low-grade intermediate-risk non-muscle invasive bladder cancer (LG-IR-NMIBC)** is back, you likely have questions. It's a lot to take in.

Use this guide to talk to your doctor about **ZUSDURI (Zuss-dur-ee)** and find out if it's right for you. Inside you'll find questions to consider and space for notes to review later.

\*Compensated for his time.

## APPROVED USE FOR ZUSDURI

ZUSDURI (mitomycin) for intravesical solution is a prescription medicine used to treat adults with a type of cancer of the lining of the bladder called low-grade intermediate-risk non-muscle invasive bladder cancer (LG-IR-NMIBC) after you have previously received bladder surgery to remove tumor and it did not work or is no longer working.

## IMPORTANT SAFETY INFORMATION

**You should not receive ZUSDURI if you** have a hole or tear (perforation) of your bladder or if you have had an allergic reaction to mitomycin or to any of the ingredients in ZUSDURI.

Please click here for [Full Prescribing Information](#), including the [Patient Information](#).




Julio is a  
ZUSDURI patient.\*

# ZUSDURI—the 1st and only FDA-approved, non-surgical treatment for recurrent LG-IR-NMIBC


## Delivers powerful, lasting results

In the ZUSDURI clinical trial, 223 patients with recurrent LG-IR-NMIBC received ZUSDURI once a week for 6 weeks. The results are as follows:



of patients  
**had no detectable tumor at 3 months**  
after treatment with ZUSDURI

(No detectable tumor is also called a complete response or CR.)



of patients who had no detectable  
tumor at 3 months were  
***still* tumor-free 12 months later**

## IMPORTANT SAFETY INFORMATION (cont'd)

**The most common side effects of ZUSDURI include:** increased blood creatinine levels, increased blood potassium levels, trouble with urination, decreased red blood cell counts, increase in certain blood liver tests, increased or decreased white blood cell counts, urinary tract infection, blood in your urine.

**Before receiving ZUSDURI, tell your healthcare provider about all of your medical conditions, including if you** have kidney problems, are pregnant or plan to become pregnant. ZUSDURI can harm your unborn baby. You should not become pregnant during treatment with ZUSDURI. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZUSDURI.

**See additional Important Safety Information throughout.**

FDA=Food and Drug Administration.

Recurrent means the cancer came back.

**2 Please click here for Full Prescribing Information,  
including the Patient Information.**

  
**Zusduri**  
(mitomycin) for intravesical solution

## How ZUSDURI is given



### 6 doses in total

- Given once a week for 6 weeks in the doctor's office or an outpatient clinic
- It's important that you receive all 6 doses of ZUSDURI



### Delivered directly in the bladder

- Your healthcare provider will give you ZUSDURI through a thin, flexible tube called a urinary catheter. The catheter will be inserted through your urethra and into your bladder (called intravesical therapy or IVT)



### Innovative gel technology

- The gel works in your bladder, where it's needed, releasing mitomycin (MY-toh-MY-sin). Mitomycin has been used extensively to treat certain cancers for many years



### There's no need for general anesthesia or surgery and patients go home the same day

## Follow all of the instructions

from your healthcare provider prior to and after treatment

## IMPORTANT SAFETY INFORMATION (cont'd)

**Females who are able to become pregnant:** You should use effective birth control (contraception) during treatment with ZUSDURI and for 6 months after the last dose.

**Males being treated with ZUSDURI:** You should use effective birth control (contraception) during treatment with ZUSDURI and for 3 months after the last dose.

Tell your healthcare provider if you are breastfeeding or plan to breastfeed. It is not known if ZUSDURI passes into your breast milk. Do not breastfeed during treatment with ZUSDURI and for 1 week after the last dose.

**Please click here for Full Prescribing Information,  
including the Patient Information.**



How is recurrent LG-IR-NMIBC different from other types of bladder cancer?

---

---

---

---

---

---

---

---

---

---

What might have caused the cancer to return?

---

---

---

---

---

---

---

---

---

---

Here are more questions to ask your doctor to see if ZUSDURI is the right treatment for you.

**1. What is ZUSDURI and how does it differ from other treatment options?**

---

---

---

---

---

**2. How well has ZUSDURI worked for people with my type of cancer?**

---

---

---

---

---

**3. Would you consider me a candidate for ZUSDURI?**

---

---

---

---

---

#### 4. How can we tell if the treatment is working?

---

---

---

---

---

#### 5. How and where is ZUSDURI given?

---

---

---

---

---

#### 6. How long will each treatment appointment last?

---

---

---

---

---

Please click here for [Full Prescribing Information](#),  
including the [Patient Information](#).

### 7. How often will I need to receive ZUSDURI?

---

---

---

---

---

### 8. What can I expect when I start treatment with ZUSDURI?

---

---

---

---

---

### 9. What are the possible side effects of ZUSDURI treatment and how long do they last? Any tips on how to manage them?

---

---

---

---

---

**10. Are there any financial support programs available for ZUSDURI?**

---

---

---

**11. What are some other things I can do to try to help reduce the risk of cancer coming back?**

---

---

---

---

---

**12. Additional questions/notes:**

---

---

---

---

Please click here for [Full Prescribing Information](#),  
including the [Patient Information](#).



# IMPORTANT SAFETY INFORMATION

**You should not receive ZUSDURI if you** have a hole or tear (perforation) of your bladder or if you have had an allergic reaction to mitomycin or to any of the ingredients in ZUSDURI.

**Before receiving ZUSDURI, tell your healthcare provider about all of your medical conditions, including if you:**

- have kidney problems
- are pregnant or plan to become pregnant. ZUSDURI can harm your unborn baby. You should not become pregnant during treatment with ZUSDURI. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZUSDURI.

**Females who are able to become pregnant:** You should use effective birth control (contraception) during treatment with ZUSDURI and for 6 months after the last dose.

**Males being treated with ZUSDURI:** You should use effective birth control (contraception) during treatment with ZUSDURI and for 3 months after the last dose.

- are breastfeeding or plan to breastfeed. It is not known if ZUSDURI passes into your breast milk. Do not breastfeed during treatment with ZUSDURI and for 1 week after the last dose.

Open discussions with your healthcare team are key.  
They want to help you make informed decisions

# IMPORTANT SAFETY INFORMATION (cont'd)

## How will I receive ZUSDURI?

- You will receive your ZUSDURI dose from your healthcare provider 1 time a week for 6 weeks into your bladder through a tube called a urinary catheter. It is important that you receive all 6 doses of ZUSDURI according to your healthcare provider's instructions.
- If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment.
- During treatment with ZUSDURI, your healthcare provider may tell you to take additional medicines or change how you take your current medicines.

## After receiving ZUSDURI:

- ZUSDURI may cause your urine color to change to a violet to blue color. Avoid contact between your skin and urine for at least 24 hours.
- To urinate, **males and females should sit** on a toilet and flush the toilet several times after you use it. After going to the bathroom, wash your hands, your inner thighs, and genital area well with soap and water.
- Clothing that comes in contact with urine should be washed right away and washed separately from other clothing.

**The most common side effects of ZUSDURI include:** increased blood creatinine levels, increased blood potassium levels, trouble with urination, decreased red blood cell counts, increase in certain blood liver tests, increased or decreased white blood cell counts, urinary tract infection, blood in your urine.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. You may also report side effects to UroGen Pharma at 1-855-987-6436.

**Please see the ZUSDURI [Full Prescribing Information](#), including the [Patient Information](#), for additional information.**



ZUSDURI™ is a trademark and UroGen® is a registered trademark of UroGen Pharma, Ltd.  
©2025 UroGen Pharma, Inc. All rights reserved.  
US-ZUS-00033 06/25

